



Anal Fissure

What is an anal fissure?

An anal fissure is a small tear or cut in the skin that lines the anus. Fissures typically cause pain and often bleed. Fissures are quite common, but are often confused with other causes of pain and bleeding, such as hemorrhoids.

A simple acute fissure, transanal view.



What are the symptoms of an anal fissure?

The typical symptoms of an anal fissure are pain during or after defecation and bleeding. Patients may try to avoid defecation because of the pain.

What causes an anal fissure?

Trauma: anything that can cut or irritate the inner lining of the anus can cause a fissure. A hard, dry bowel movement is typically responsible for a fissure. Other causes of a fissure include diarrhea or inflammatory conditions of the anal area. Anal fissures may be acute (recent onset) or chronic (present for a long time or recurring frequently). Chronic fissures often have a small external lump associated with the tear called a sentinel pile or skin tag.

How can a fissure be treated?

Often treating one's constipation or diarrhea can cure a fissure. An acute fissure is typically managed with non-operative treatments and over 90% will heal without surgery. A high fiber diet, bulking agents (fiber supplements), stool softeners, and plenty of fluids help relieve constipation, promote soft bowel movements, and aide in the healing process. Increased dietary fiber may also help to improve diarrhea. Warm baths for 10-20 minutes several times each day are soothing and promote relaxation of the anal muscles, which can also help healing. Occasionally, special medications may be recommended. A chronic fissure may require additional treatment.

Will the problem return?

Fissures can recur easily, and it is quite common for a healed fissure to recur after a hard bowel movement. Even after the pain and bleeding has disappeared one should continue to aim for good bowel habits and adhere to a high fiber diet or fiber supplement regimen. If the problem returns without an obvious cause, further assessment may be needed.

What can be done if a fissure doesn't heal?


A fissure that fails to respond to treatment should be re-examined. Persistent hard or loose bowel movements, scarring, or spasm of the internal anal sphincter muscle all contribute to delayed healing. Other medical problems such as inflammatory bowel disease, infections, or anal growths (skin tumors) can cause fissure-like symptoms, and patients suffering from persistent anal pain should be examined to exclude these conditions.

What does surgery involve?

Surgery is a highly effective treatment for a fissure and recurrence rates after surgery are low. Surgery usually consists of a small operation to cut a portion of the internal anal sphincter muscle (a lateral internal sphincterotomy). This helps the fissure heal and decreases pain and spasm.

If a sentinel pile is present, it too may be removed to promote healing of the fissure. A sphincterotomy rarely interferes with one's ability to control bowel movements and is most commonly performed as a short outpatient procedure. Surgical treatments do have other risks, and your surgeon will address these with you.

A chronic fissure with a skin tag and a hypertrophied anal papilla, coronal view.



How long does the healing process take after surgery?

Complete healing occurs in a few weeks, although pain often disappears after a few days.

Can fissures lead to colon cancer?

No! Persistent symptoms, however, need careful evaluation since conditions other than fissure can cause similar symptoms. Your doctor may request additional testing even if your fissure has successfully healed. A colonoscopy may be required to exclude other causes of bleeding.

© 2008 American Society of Colon & Rectal Surgeons

R10/06 - P01/08